

## **Cambridge Assessment International Education**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

BIOLOGY 0610/63

Paper 6 Alternative to Practical

October/November 2019

1 hour

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



© UCLES 2019

1 Photosynthesis occurs in chloroplasts. A chloroplast suspension can be made by extracting the chloroplasts from green leaves.

The blue dye DCPIP can be used to investigate photosynthesis. During photosynthesis the blue colour of the DCPIP changes and eventually disappears so that it becomes colourless. The remaining mixture appears green due to the chloroplast suspension.

A student investigated the effect of light on photosynthesis in a chloroplast suspension.

- The student crushed eight green leaves in an ice-cold buffer solution. Step 1
- Step 2 The mixture of crushed leaves and buffer solution was then filtered to form a chloroplast suspension.
- Step 3 The student was given two test-tubes. One test-tube was wrapped in aluminium foil to exclude light. The second test-tube was not covered in foil so that light could reach the contents of this test-tube.
- A plastic pipette was used to add 2 cm<sup>3</sup> of the chloroplast suspension to each of the two Step 4 test-tubes.
- A syringe was used to add 5 cm<sup>3</sup> of DCPIP solution to both test-tubes. A stopper was Step 5 placed in each test-tube.
- Step 6 Both test-tubes were placed in a water-bath containing ice-cold water. The water-bath was positioned near a bright light.
- Step 7 The test-tubes were left in the bright light for five minutes.
- Step 8 After five minutes the student observed and noted the colour of the liquid in the test-tube that was **not** wrapped in foil. They removed the foil from the covered test-tube and observed the colour in this test-tube.

The student's results are shown in Fig. 1.1.




Fig. 1.1

© UCLES 2019

(a) (i) Prepare a table and record the results from Fig. 1.1.

	[2
(ii)	Identify the variable that was changed (independent variable) and the variable that was measured (dependent variable) in this investigation.
	independent variable
	dependent variable
	[2
(iii)	State <b>two</b> variables that were kept constant in this investigation.  1
	2[2
(iv)	State a conclusion for these results.
	[1

**(b)** In step 6 both test-tubes were placed in a water-bath containing ice-cold water for five minutes. A thermometer was also placed in the water-bath.

Fig. 1.2 shows the thermometer at the start of the investigation and after five minutes.




Fig. 1.2

Record the temperatures shown in Fig. 1.2. Include the units.
temperature at the start of the investigation
temperature after five minutes
[2
The investigation was not repeated.
(i) Suggest why it would be advisable to repeat the investigation.

© UCLES 2019 0610/63/O/N/19

(c)

	(ii)	Identify <b>one</b> other source of error in this investigation and suggest an improvement for this error.
		error
		improvement
(d)		[2] tudent wanted to investigate the effect of temperature on the rate of photosynthesis in a proplast suspension.
	Des	scribe how the student could carry out this investigation using DCPIP.
		[6]
		FT 1 400

[Total: 18]

**2** Fig. 2.1 is a photomicrograph of part of the lower epidermis of a leaf.




Fig. 2.1

(a) (i) Make a large drawing of Fig. 2.1. Do not label your drawing.

[4]

	(ii)	Measur	e the le	ength o	of the li	ine MN	l on Fi	g. 2.1.					
		length o	of line <b>I</b>	ЛN					mm				
		Calcula Fig. 2.1									ndicat	ed by li	ne <b>MN</b> in
				magn	ificatio	n = <del>act</del>	le ual wic	ength of t	of line <b>I</b> he gua	MN on ird cell:	Fig. 2.	1 the stor	<del></del>
		Give yo	ur ans						J				
													mm
													[3]
(b)	A st	tudent in	vestiga	ted the	effect	t of suc	crose c	oncen	tration	on sto	matal	opening	<b>J</b> .
	Stri	ps of epi	dermis	from le	eaves	were p	laced i	n diffe	rent co	ncentra	ations	of sucr	ose solution for
		hour.				·							
		student ution.	meası	ired th	e width	of 10	stoma	tal ope	enings	from th	e epid	ermis i	n each sucrose
	The	results a	are sho	wn in	Table 2	2.1.							
						Т	able 2	.1					
nor	oont	222							atal an	onina /			
•	ercentage centration of sucrose			_	0				· ·	ening /	•	40	
SI			1	2	3	4	5	6	7	8	9	10	average
	0 5		5 3	6	7	5 3	5	6	7	5 3	7 3	6	5.9
	20		1	2	1	2	1	1	5	1	2	1	1.7
	20		<u>'</u>		•		<u> </u>	•		'		<u> </u>	1.7
	(i)	Circle o	<b>ne</b> me	asurer	nent in	Table	2.1 tha	at coul	d be ar	nomalo	us.		[1]
	(ii)	Calcula	te the a	averag	e widtl	n of sto	matal	openir	ng in th	e 5% s	sucros	e soluti	on.
		Include	the un	it.									
													[2]

(c) Pineapple plants are adapted to grow in hot dry conditions.

Scientists investigated the number of stomata open, in pineapple plants, at different times of day.

The data is shown in Table 2.2 (00:00 is midnight).

Table 2.2


time of day	average number of stomata open per mm <sup>2</sup>
00:00	77
04:00	61
08:00	22
12:00	4
16:00	10
20:00	51

(i)	Calculate the percentage change in the average number of open stomata per mr	$n^2$
	petween 16:00 hours and 20:00 hours.	

Space for working.

 	%
	[2]

(ii) Plot a bar chart on the grid to show the data in Table 2.2.



(iii)	Describe, using your graph, how the average number of open stomata per mm <sup>2</sup> chan throughout the day.	ges

[3]

Wh	en fruit ripens the starch contained within it is converted into reducing sugars.
(i)	State the name of the substance that is used to test for the presence of starch.
	[1]
(ii)	Describe how you could safely show a reducing sugar was present in a sample of pineapple fruit juice.
	procedure
	safety
	[4]
	[Total: 22]

11

## **BLANK PAGE**

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.