



Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

BIOLOGY

0610/52

Paper 5 Practical Test May/June 2024

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].

For Examiner's Use						
1						
2						
Total						

This document has 12 pages. Any blank pages are indicated.





1 You are going to investigate the effect of acid concentration on the rate of diffusion.

Bromothymol blue is an indicator that turns yellow in the presence of an acid.

Read all the instructions but DO NOT DO THEM until you have drawn a table for your results in the space provided in 1(a)(iii).

2

You should use the safety equipment provided while you are doing the practical work.

- Step 1 Label three beakers A1, A2 and A3.
- Step 2 Add 20 cm³ of 1 mol per dm³ hydrochloric acid solution to beaker **A1**.
- Step 3 Add 2cm³ of 1 mol per dm³ hydrochloric acid solution to beaker **A2**.
- Step 4 Add 18 cm³ of distilled water to the hydrochloric acid solution in beaker **A2** and mix well.
- Step 5 Use a clean syringe to transfer 2 cm³ of the solution from beaker **A2** to beaker **A3**.
- Step 6 Add 18 cm³ of distilled water to beaker **A3** and mix well.

Table 1.1 shows the final concentrations of the hydrochloric acid solution in each beaker.

Table 1.1

beaker	final concentration of the hydrochloric acid solution / mol per dm ³	final volume of the solution/cm ³
A 1	1.00	20
A2	0.10	
А3	0.01	20

(a) (i) Complete Table 1.1 by calculating the final volume of the solution in beaker A2.

Write your answer in Table 1.1.

[1]

You are provided with three test-tubes containing bromothymol blue indicator in agar jelly.

- Step 7 Label the three test-tubes A1, A2 and A3.
- Step 8 Use a clean syringe to transfer 5 cm³ of the hydrochloric acid solution from beaker **A1** to test-tube **A1**.
- Step 9 Use a clean syringe to transfer 5 cm³ of the hydrochloric acid solution from beaker **A2** to test-tube **A2**.
- Step 10 Use a clean syringe to transfer 5 cm³ of the hydrochloric acid solution from beaker **A3** to test-tube **A3**.



* 0019655390503 *

3

- Step 11 Raise your hand when you are ready for warm water to be poured into the beaker labelled **water-bath**.
- Step 12 Place the three test-tubes into the warm water-bath and start the stop-clock.

Leave the test-tubes in the water-bath for 20 minutes.

While you are waiting continue with the other questions.

- Step 13 After 20 minutes, remove the test-tubes from the water-bath and place them in the test-tube rack.
 - (ii) Record the temperature of the water in the water-bath.

temperature°C [1]

- Step 14 Remove the hydrochloric acid solution from each test-tube by pouring the solution in each test-tube into the beaker labelled **waste**.
- Step 15 For each test-tube, measure the distance between the top of the agar and the point at which the blue agar has become yellow. Fig. 1.1 shows where to take the measurement.

Record your measurements in your table in 1(a)(iii).




Fig. 1.1

4

(iii) Prepare a table and record your results in the space provided.

Include the final concentration of hydrochloric acid in your table.

		[4]
(iv)	State a conclusion for your results.	
		[1]
(v)	Suggest why the temperature of the water-bath did not need to be maintained do the investigation.	ıring
		[1]

* 0019655390505 * (vi) Suggest why the hydrochloric acid was removed from the test-tube in step 14 before

5

` '	measuring the distance of the yellow colour.	
		. [1]
(vii)	Identify the independent variable and the dependent variable in this investigation.	
	independent variable	
	dependent variable	
		[2]
(viii)	Explain why suitable eye protection should be worn when doing this investigation.	
		[4]

(b) In an experiment, a student cut different-sized cubes of agar containing bromothymol blue indicator. The student placed the cubes in hydrochloric acid and measured the time taken for each cube to become yellow.

6

The results are shown in Table 1.2.

Table 1.2

length of the cube sides/mm	volume of the cube/mm ³	time taken for the cubes to become yellow/s				
2	8	35				
4		60				
6	216	170				
8	512	275				
10	1000	445				


(i) Complete Table 1.2 by calculating the volume of the 4 mm cube.

Write your answer in Table 1.2.

[1]

* 0019655390607 *

(ii) Plot a line graph on the grid of the length of the cube sides against the time taken for the



7

(iii)	Use your graph to estimate the time taken for a cube with a side length of 3mm to become yellow.
	Show on your graph how you obtained your answer.
	s [2]
(iv)	State two variables that should have been kept constant in this investigation.
	1
	2
	[2]

[Total: 21]

[4]

(a) When a seed germinates it grows into a seedling.

Fig. 2.1 is a photograph of a seedling.



8

Fig. 2.1

(i) Make a large drawing of the seedling shown in Fig. 2.1.



9

(ii) Line AB represents the width of the seedling.

Measure the length of line AB on Fig. 2.1.

length of line ABmm


The actual width of the seedling is 9mm.

Calculate the magnification of the seedling in Fig. 2.1 using the formula and your measurement.


magnification =
$$\frac{\text{length of line } \mathbf{AB} \text{ in Fig. 2.1}}{\text{actual width of the seedling}}$$

Give your answer to two significant figures.

Space for working.



(iii) Fig. 2.2 shows photographs of two different seedlings, P and Q.



10

Fig. 2.2

Describe one visible similarity and two visible differences between seedling ${\bf P}$ and

	seedling Q in Fig. 2.2.	
	similarity 1	
	difference 1	
	difference 2	
		 [3]
		[၁]
(b)	As seedlings grow, they change starch into reducing sugars.	
	Describe the methods you would use to test a sample of seedlings for starch a reducing sugars.	and
	starch test	
	reducing sugar test	



(c) When light shines from one side, seedlings grow towards the light.

Fig. 2.3 shows a diagram of a seedling growing towards a light source. The angle of growth is shown on the diagram.

11




Fig. 2.3

Plan an seedling	gation	to fin	d out	how	light	intensity	affects	the	angle	at	which	the
	 											[6]

[Total: 19]

12

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

