

Cambridge IGCSE[™]

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

CHEMISTRY 0620/33

Paper 3 Theory (Core)

October/November 2021

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

1 (a) A list of formulae is shown.


Br₂
CO
CO₂
CH₄
C₂H₆
HC*l*KBr
LiC*l*MgC*l*₂
O₂

Answer the following questions using these formulae. Each formula may be used once, more than once or not at all.

State which formula represents:

(1)	a compound that gives a red colour in a flame test	
		[1]
(ii)	a compound containing an ion with a 2+ charge	
		[1]
(iii)	a compound that is a product of respiration	
		[1]
(iv)	a compound used as a food preservative	
		[1]
(v)	an element that is used in the production of steel.	
		[1]

(b) Complete the dot-and-cross diagram to show the electron arrangement in a molecule of hydrogen chloride. Show outer shell electrons only.



[2]

(c)	State whether carbon dioxide is a basic oxide or an acidic oxide. Give a reason for your answer.
	[1]
	[Total: 8]

2 The table shows the masses of some of the ions in 1000 cm³ of water from a river.

(a) Answer these questions using only the information in the table.

name of ion	formula of ion	mass of ion in 1000 cm ³ of river water/mg
ammonium	NH ₄ ⁺	1.0
	Ca ²⁺	16.5
chloride	Cl ⁻	7.0
iron(III)	Fe³+	0.5
magnesium	Mg ²⁺	4.0
	NO ₃ -	0.5
potassium	K ⁺	3.5
silicate	SiO ₃ ²⁻	7.5
sodium	Na⁺	6.0
sulfate	SO ₄ ²⁻	11.0

	(i)	State which positive ion has the lowest concentration.
		[1]
	(ii)	Name the compound containing Ca ²⁺ and NO ₃ ⁻ ions. [1]
(iii)	Calculate the mass of sulfate ions in 500 cm ³ of river water.
		mass = mg [1]
(b)	Des	scribe a test for iron(III) ions.
	test	
	obs	ervations

[2]

(c) Compound **A** is found in river water. The structure of compound **A** is shown.

(i) On the structure draw a circle around the alcohol functional group. [1]

(ii) Deduce the formula of compound **A** to show the number of carbon, hydrogen, oxygen and nitrogen atoms.

.....[1]

(iii) Another compound found in river water has the formula C₄H₈O₂.

Complete the table to calculate the relative molecular mass of this compound.

type of atom	number of atoms	relative atomic mass	
carbon	4	12	4 × 12 = 48
hydrogen		1	
oxygen		16	

relative molecular mass =[2]

[Total: 9]

	is extracted in a blast furnace using a mixture of iron ore, coke (carbon), air and calcium carbon estone).	ate
(a)	Give two reasons why air is blown into the blast furnace.	
	1	
	2	[2]
(b)	Magnetite is an ore of iron which contains a compound of iron with the formula $\mathrm{Fe_3O_4}$.	
	(i) Give the name of another ore of iron.	
		[1]
((ii) In the blast furnace Fe ₃ O ₄ is reduced to Fe.	
	Complete the chemical equation for the reduction of Fe ₃ O ₄ .	
	$Fe_3O_4 + 4CO \rightarrowFe +CO_2$	[2]
(i	iii) Explain how this equation shows that Fe ₃ O ₄ is reduced.	
		[1]
	Calcium carbonate (limestone) is added to the blast furnace. The calcium carbonate undergoes thermal decomposition.	
	Give the meaning of the term thermal decomposition.	
		[2]
(d)	Iron can form alloys such as vanadium steel.	
	(i) State the meaning of the term <i>alloy</i> .	
		[1]

(ii)	Choose from the	diagrams, B , C , D or	E , the structure wh	ich best represents an allo	y.
	В	С	D	E	
			5	structure	[1]
(iii)	The symbol for a	an isotope of vanadiun	n is shown.		
		51 ¹ 23	V		
	Deduce the nunvanadium.	nber of electrons, neu	utrons and protons	in one atom of this isoto	pe of
	number of electr	ons			
	number of neutro	ons			
	number of proto	ns			
					[3]
(iv)	Vanadium is ma	lleable and conducts e	electricity.		
	Give two other p	physical properties of	vanadium that are o	haracteristic of all metals.	
	1				
	2				
					[2]

[Total: 15]

4 The table shows some properties of four halogens in Group VII.

halogen	melting point /°C	boiling point /°C	density of liquid at boiling point in g/cm³
chlorine	-101	-35	1.56
bromine	-7	59	
iodine	114		4.93
astatine	302	337	6.35

	(a)	ı (i) Com	nlete	the	table	hv	predictin	a	ŀ
١	a	, (1		picic	uic	labic	υy	predictiri	ч	

•	the boiling point of iodine
•	the density of bromine.

[2]

(II)	Describe the trend in	the melting points o	of the halogens down	Group VII.

	[1]

(b) (i) Deduce the electronic structure of chlorine. Use the Periodic Table to help you.


[1	[[
----	----

(ii) Explain why a bromide ion has a single negative charge.

 	 [1]

(c) Scientists have predicted that sodium astatide reacts with chlorine.

Complete the word equation for this reaction.



[2]

(d)	Hydrogen chloride is produced when chlorine reacts with ammonia. An aqueous solution of hydrogen chloride is acidic.
	Describe how you could determine the pH of an acidic solution without using a pH meter.
	[2]
	[Total: 9

5 The table shows the structures of some organic compounds.

compound	structure of compound	homologous series
F	H H O H—C—C—C—O—H H H	carboxylic acid
G	H H	
Н	H H H H—C—C—C—H H H H	

(a) Complete the table by naming the homologous series.

The first one has been done for you.


[2]

(b) Draw the structure of a compound containing two carbon atoms which belongs to the same homologous series as compound F. Show all of the atoms and all of the bonds.

[1]

(c)	Com	pound	G	reacts	with	bromine
-----	-----	-------	---	--------	------	---------

Choose from the structures, P, Q, R or S, the structure of the product formed.



F 4	4.7
17	
	4

(d) (i) Compound G can be obtained by cracking petroleum fractions.	(d)	(i)	Compound	G c	can be	obtained	by	cracking	petroleum	fractions.
--	-----	-----	----------	------------	--------	----------	----	----------	-----------	------------

Describe what is meant by the term *cracking*.

- (ii) Name the product of the reaction when compound **G** reacts with steam.
- (iii) Many molecules of compound **G** can join together to form a compound with a very long chain.

Choose from the list the general name given to a compound that is formed by the addition of many small units.

......[1]

Draw a circle around the correct answer.


isomer monomer naphtha polymer [1]


(e) State the names of the two compounds formed during the complete combustion of compound H.


..... and [2]

[Total: 10]

6 The diagram shows part of the structures of sodium chloride and pentane at room temperature and pressure.







1	a١	Describe the	differences in	the physical	properties of these	substances in terms	of
ı	a)	Describe the	differences in	the physical	properties of these	substances in terms	5 OI.

•	volatility

sodium chloride	
pentane	
solubility in water	
sodium chloride	
pentane	
electrical conductivity when molten (liquid).	
sodium chloride	
pentane	
	[5]

(b) Concentrated aqueous sodium chloride is electrolysed using carbon (graphite) electrodes.
Name the gas produced at the positive electrode.

______[1]

- (c) Pentane is a fuel. Under some conditions pentane forms carbon monoxide.
 - (i) State the condition under which pentane forms carbon monoxide.

 [1]
 - (ii) State one adverse effect of carbon monoxide on health.

.....[1]

[Total: 8]

7

(i)	Name the change of state from solid to liquid.	[1]
(ii)	Use the kinetic particle theory to describe the differences between solid aluminium liquid aluminium in terms of:	
	the separation of the particles	
	the motion of the particles.	
		[4]
(b) Alu	uminium is extracted from aluminium ore by electrolysis.	
		[4]
Ex	uminium is extracted from aluminium ore by electrolysis. plain why aluminium is extracted by electrolysis and not by reduction with carbon.	[4]
(c) Giv	uminium is extracted from aluminium ore by electrolysis. plain why aluminium is extracted by electrolysis and not by reduction with carbon. We two reasons why aluminium is used in the manufacture of aircraft.	[4]
(c) Giv 1.	uminium is extracted from aluminium ore by electrolysis. plain why aluminium is extracted by electrolysis and not by reduction with carbon. We two reasons why aluminium is used in the manufacture of aircraft.	[4]

(e) The table compares the ease of reduction of four metal oxides when heated with carbon.

metal oxide	ease of reduction					
aluminium oxide	not reduced at 2080 °C					
nickel(II) oxide	reduced at 540 °C					
titanium(IV) oxide	reduced at 1600°C					
zinc oxide	reduced at 850°C					

Put the four metals in order of their reactivity. Put the least reactive metal first.


	leas	st reactive most reactive	[2]
(f)	Met	thane is used as a fuel in the extraction of some metals.	
	(i)	State the main source of methane.	
			[1]
	(ii)	State one adverse effect of methane on the environment.	
			[1]
		[Total:	13]

8 A student investigated the reaction of small pieces of zinc with dilute hydrochloric acid. The hydrochloric acid was in excess.

$$Zn + 2HCl \rightarrow ZnCl_2 + H_2$$

The rate of reaction is found by measuring the increase in volume of hydrogen gas with time.

The results are shown on the graph.



(a) Deduce the time taken from the beginning of the experiment to collect 30 cm³ of hydrogen gas.

(b) The experiment was repeated using dilute hydrochloric acid of a higher concentration.

All other conditions stayed the same.

Draw a line **on the grid** to show how the volume of hydrogen gas changes with time. [2]

(c)	Describe the effect each of the following has on the rate of reaction of zinc with dilute hydrochloric acid.
	All other conditions stay the same.
	The reaction is carried out at a lower temperature.
	The reaction is carried out using zinc powder instead of small pieces of zinc.
	[2]
(d)	When 0.065 g of zinc is used, 24 cm³ of hydrogen gas is formed. Calculate the mass of zinc needed to produce 96 cm³ of hydrogen gas.
	mass of zinc = g [1]
(e)	Aqueous ammonia is added to aqueous zinc chloride.
	Describe the observations when a few drops of aqueous ammonia are added and when excess aqueous ammonia is added.
	a few drops of aqueous ammonia
	excess aqueous ammonia
	[2]
	[Total: 8

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

		2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon			
	II/			6	Щ	fluorine 19	17	Cl	chlorine 35.5	35	ğ	bromine 80	53	П	iodine 127	85	¥	astatine -			
	IA			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>L</u>	tellurium 128	84	Ъ	moloum –	116		livermorium -
	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209			
	2			9	ပ	carbon 12	14	:S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
	≡			2	В	boron 11	13	Ν	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> 1	thallium 204			
										30	Zu	zinc 65	48	S	cadmium 112	80	Нg	mercury 201	112	ü	copernicium -
										29	Cn	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium -
Group										28	Ż	nickel 59	46	Pd	palladium 106	78	చ	platinum 195	110	Ds	darmstadtium -
Ď										27	ဝိ	cobalt 59	45	格	rhodium 103	77	'n	iridium 192	109	Ĭ	meitnerium -
		- I	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium -
							1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
				_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	>	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	Б	tantalum 181	105	o D	dubnium -
					atc	rel				22	j	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	¥	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	56	Ba	barium 137	88	Ra	radium
	_			8	=	lithium 7	7	Na	sodium 23	19	×	potassium 39	37	&	rubidium 85	22	S	caesium 133	87	Ļ	francium -

71 Lu	lutetium 175	103	۲	lawrencium	ı
°° Yb	ytterbium 173	102	8	nobelium	ı
mL Tm	thulium 169	101	Md	mendelevium	ı
es Fr	erbium 167	100	Fm	fermium	I
67 Ho	holmium 165	66	Es	einsteinium	ı
66 Dy	dysprosium 163	86	ర్	californium	I
65 Tb	terbium 159	97	BK	berkelium	ı
Gd Gd	gadolinium 157	96	Cm	curium	ı
63 Eu	europium 152	92	Am	americium	I
62 Sm	samarium 150	94	Pn	plutonium	ı
61 Pm	promethium -	93	ď	neptunium	ı
9 9 8	neodymium 144	92	\supset	uranium	238
59 Pr	praseodymium 141	91	Ра	protactinium	231
Se Ce	cerium 140	06	드	thorium	232
57 La	lanthanum 139	89	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).