



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



CHEMISTRY

0620/42

Paper 4 Theory (Extended)

May/June 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has **16** pages. Any blank pages are indicated.



- 1 A list of gases is shown.

ammonia
helium
hydrogen
carbon dioxide
carbon monoxide
chlorine
methane
nitrogen dioxide
propene
sulfur dioxide

Answer the following questions about these gases.

Each gas may be used once, more than once or not at all.

- (a) State **one** gas which:

- (i) is the main constituent of natural gas

..... [1]

- (ii) is responsible for both photochemical smog **and** acid rain

..... [1]

- (iii) is unsaturated

..... [1]

- (iv) has monatomic particles

..... [1]

- (v) reduces iron(III) oxide in a blast furnace.

..... [1]

- (b) Nitrogen dioxide, NO_2 , and carbon monoxide are removed from a car exhaust by a catalytic converter.

Write the symbol equation for this reaction.

..... [2]

[Total: 7]





- 2 A list of five metals is shown.

copper
iron
magnesium
potassium
silver

- (a) All metals form positive ions.

- (i) Describe how atoms form positive ions.

..... [1]

- (ii) State which of the five metals in the list has the greatest tendency to form positive ions.

..... [1]

- (iii) Suggest **one** of the five metals in the list which is **not** likely to show catalytic properties.

..... [1]

- (iv) State which of the five metals in the list is a major component of stainless steel.

..... [1]

- (b) A student adds a sample of a metal to an aqueous metal salt in a beaker to see if a displacement reaction takes place.

Complete Table 2.1 to show the colour of the solution in the beaker at the start and at the end of the experiment.

Table 2.1

metal	aqueous solution	colour at the start	colour at the end
magnesium	iron(II) sulfate	green	
silver	copper(II) sulfate		

[3]





- (c) Most Group II metals form a gas when placed into cold water. An alkaline solution is also formed.

(i) Name the gas formed when strontium is added to cold water.

..... [1]

(ii) Name the alkaline solution formed when strontium is added to cold water.

..... [1]

- (iii) One Group II metal reacts very slowly when placed in cold water. When heated, the metal reacts with steam to form a white solid.

Identify this metal and name the white solid formed.

metal

white solid

[2]

- (d) Under certain conditions, iron will react with steam to form an oxide of iron with the formula Fe_3O_4 .

Fe_3O_4 reacts with dilute hydrochloric acid to form a mixture of iron(II) and iron(III) salts and water.

Deduce the symbol equation for the reaction between Fe_3O_4 and dilute hydrochloric acid.


..... [3]

[Total: 14]





- DO NOT WRITE IN THIS MARGIN
- 3 The symbol equation for the industrial production of ammonia is shown.



- (a) Name this industrial process.

..... [1]

- (b) State the meaning of ΔH .

..... [1]

- (c) State the typical conditions and name the catalyst used in the industrial production of ammonia.

temperature and units

pressure and units

catalyst used

[3]

- (d) State **two** methods of increasing the rate of this reaction.

1

2

[2]





(e) The symbol equation for the reaction can be represented as shown in Fig. 3.1.




Fig. 3.1

Table 3.1 shows some bond energies.

Table 3.1

bond	N≡N	H–H
bond energy in kJ/mol	945	435

Use the bond energies in Table 3.1 and ΔH to calculate the bond energy of an N–H bond, in kJ/mol.

Use the following steps.

- Calculate the energy needed to break bonds in the reactants.

..... kJ

- Calculate the energy released when bonds form in the products.

..... kJ


- Calculate the energy of an N–H bond.

..... kJ/mol
[3]





(f) An incomplete symbol equation for the preparation of ammonia in the laboratory is shown.



(i) Complete the symbol equation.

[1]

(ii) Name NH_4Cl .

..... [1]

(iii) Calculate the volume of ammonia, NH_3 , measured at room temperature and pressure, which forms when 1.12g of CaO is heated with excess NH_4Cl .
[M_r: CaO, 56]

..... cm³ [3]

[Total: 15]





- 4 A carboxylic acid reacts with an alcohol to produce an ester and water.

Under certain conditions, this reaction can be reversed so an ester reacts with water to produce a carboxylic acid **X** and an alcohol **Y**.

The reaction reaches an equilibrium.



The forward reaction is endothermic.

- (a) Deduce the empirical formula of the ester.

..... [1]

- (b) Name the ester.

..... [1]

- (c) Name carboxylic acid **X** and draw its displayed formula.

name

displayed formula

[2]

- (d) Name alcohol **Y** and give its structural formula.

name

structural formula

[2]





- (e) Complete Table 4.1 to show the effect, if any, for each change of condition.

Table 4.1

change of condition	effect on the concentration of carboxylic acid X at equilibrium
temperature is decreased	
concentration of $\text{CH}_3\text{CH}_2\text{COOCH}_3$ is decreased	
more alcohol Y is added	
a catalyst is added	

[4]

- (f) At the beginning of the reaction between the ester and water, no carboxylic acid is present in the reaction mixture.

- (i) Suggest how the pH of the reaction mixture changes from the start of the reaction until equilibrium is reached.

Assume alcohols and esters are neutral.

pH at start of reaction

pH at equilibrium

[2]

- (ii) Identify the ion that causes the change in pH.

..... [1]

- (iii) Name an indicator which can be used to follow the change in pH.

..... [1]

[Total: 14]





5 Sulfur is a Group VI element.

(a) A sample of sulfur contains two isotopes, ^{32}S and ^{34}S .

(i) Complete Table 5.1 to show the number of protons and neutrons in one atom of each isotope of sulfur.

Table 5.1

	^{32}S	^{34}S
protons		
neutrons		

[2]

(ii) State why these isotopes have identical chemical properties.

..... [1]

(iii) State the mass of 6.02×10^{23} atoms of ^{34}S . Include units in your answer.

..... [1]

(iv) State the name of the amount of substance which contains 6.02×10^{23} atoms.

..... [1]

(v) Table 5.2 shows the relative abundance of these isotopes of sulfur in the sample.

Table 5.2

atom	^{32}S	^{34}S
relative abundance	95%	5%

Calculate the relative atomic mass of sulfur in this sample to **one** decimal place.

relative atomic mass = [2]





(b) Sulfur reacts with magnesium to form magnesium sulfide, MgS, an ionic compound.

(i) Complete the dot-and-cross diagram in Fig. 5.1 of the ions in magnesium sulfide.

Give the charges on the ions.




Fig. 5.1

[3]

(ii) State why MgS has a high melting point.

..... [1]

(iii) State why molten MgS conducts electricity.

..... [1]

(c) An acid containing sulfur reacts with sodium hydroxide, NaOH, to form a salt and water. The salt has the formula Na₂SO₃.

(i) Deduce the formula of this acid.

..... [1]

(ii) Deduce the formula of the anion in Na₂SO₃.

..... [1]

(d) Na₂SO₃ is oxidised by acidified aqueous potassium manganate(VII).

(i) State what VII refers to in the name potassium manganate(VII).

..... [1]

(ii) State the colour change when this reaction happens.

from to [2]

[Total: 17]





6 Glucose is involved in two processes.

(a) Glucose, C₆H₁₂O₆, is made in plants from carbon dioxide and water.

(i) Name this process.

..... [1]

(ii) Write the symbol equation for this process.

..... [1]

(iii) State **two** essential conditions needed for this process to happen.

1

2

[2]

(b) Glucose is converted to ethanol.

(i) Name this process.

..... [1]

(ii) Name the **other** product formed when glucose is converted to ethanol.

..... [1]

(c) Ethanol is made by reacting ethene with steam in an industrial process.

(i) State the conditions and type of catalyst used in this industrial production of ethanol.

temperature and units

pressure and units

type of catalyst used

[3]

(ii) Explain why this reaction is an addition reaction.

..... [1]





(iii) Complete the dot-and-cross diagram in Fig. 6.1 of a molecule of ethanol.




Fig. 6.1

[3]

[Total: 13]



* 0019654852514 *



14

BLANK PAGE

DO NOT WRITE IN THIS MARGIN





BLANK PAGE

DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.





The Periodic Table of Elements

I		II		Group															
				III						IV			V		VI		VII		
Key	atomic number name relative atomic mass	atomic symbol	hydrogen 1	H															
Li lithium 7	Be beryllium 9																		
Na sodium 23	Mg magnesium 24																		
K potassium 39	Ca calcium 40	Sc scandium 45	Ti titanium 48	V vanadium 51	Cr chromium 52	Mn manganese 55	Fe iron 56	Co cobalt 59	Ni nickel 59	Cu copper 64	Zn zinc 65	Ga gallium 70	Ge germanium 73	As arsenic 75	Se selenium 79	Br bromine 80	Kr krypton 84		
Rb rubidium 85	Sr strontium 88	Y yttrium 89	Zr zirconium 91	Nb niobium 93	Tc technetium –	Mo molybdenum 96	Pd palladium 106	Rh rhodium 103	Ru ruthenium 101	Ag silver 108	Cd cadmium 112	In indium 115	Sn tin 119	Sb antimony 122	Te tellurium 128	I iodine 127	Xe xenon 131		
Cs caesium 133	Ba barium 137	La lanthanoids 56	Hf hafnium 71	Ta tantalum 72	W tungsten 73	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	Tl thallium 204	Pb lead 207	Bi bismuth 209	Po polonium –	At astatine –	Rn radon –		
Fr francium –	Ra radium –	Ac actinoids 88	Rf rutherfordium 103	Db dubnium 104	Sg seaborgium 105	Bh bohrium 106	Hs hassium 107	Mt meitnerium 108	Ds darmstadtium 109	Rg roentgenium 110	Cn copernicium 111	Nh nilssonium 112	Fl fermium 113	Mc moscovium 114	Lv livermorium –	Ts tennessine –	Og oganesson –		
lanthanoids	La lanthanum 139	Ce cerium 140	Pr praseodymium 141	Nd neodymium 144	Pm promethium –	Sm samarium 150	Eu europium 152	Gd gadolinium 157	Tb terbium 159	Dy dysprosium 163	Ho holmium 165	Er erbium 167	Tm thulium 169	Yb ytterbium 173	Lu lutetium 175				
actinoids	Ac actinium –	Th thorium 232	Pa protactinium 231	U uranium 238	Np neptunium –	Pu plutonium –	Am americium –	Cm curium –	Bk berkelium –	Cf californium –	Esn einsteinium –	Md mendelevium –	Fm fermium –	No nobelium –	Ro lawrencium –	Fr francium –			

57 La lanthanum 139	58 Ce cerium 140	59 Pr praseodymium 141	60 Nd neodymium 144	61 Pm promethium –	62 Sm samarium 150	63 Eu europium 152	64 Gd gadolinium 157	65 Tb terbium 159	66 Dy dysprosium 163	67 Ho holmium 165	68 Er erbium 167	69 Tm thulium 169	70 Yb ytterbium 173	71 Lu lutetium 175
89 Ac actinium –	90 Th thorium 232	91 Pa protactinium 231	92 U uranium 238	93 Np neptunium –	94 Pu plutonium –	95 Am americium –	96 Cm curium –	97 Bk berkelium –	98 Cf californium –	99 Es einsteinium –	100 Fm fermium –	101 Md mendelevium –	102 No nobelium –	103 Fr francium –

The volume of one mole of any gas is 24 dm^3 at room temperature and pressure (r.t.p.).

